arXiv:0803.1122v1 [math.NT] 7 Mar 2008

A NOTE ON LARSEN'S CONJECTURE AND RANKS OF ELLIPTIC CURVES

TIM[†] AND VLADIMIR DOKCHITSER

1. Introduction

In [8] M. Larsen proposed the following:

Conjecture 1. *Let K be a finitely generated infinite field, K_s its separable closure, G a finitely generated subgroup of* $\mathrm{Gal}(K_s/K)$, *and K_s^G its fixed field. If E/K is an elliptic curve, then the group $E(K_s^G)\otimes\mathbb{Q}$ is infinitely generated.*

When G is generated by one element, this conjecture is known for $K=\mathbb{Q}$ [5] and in a number of other cases [1, 6, 7]; see also [9] for some results when G has two generators. We give evidence for the full conjecture for $K=\mathbb{Q}$ by showing that it is implied both by the Shafarevich-Tate and the (first part of) the Birch–Swinnerton-Dyer conjecture for elliptic curves over number fields:

Theorem 2. *Let $E/\mathbb{Q}$ be an elliptic curve and $G\subset \mathrm{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ a finitely generated subgroup. The analytic rank and the p^∞-Selmer rank for every odd p are unbounded in number fields contained in $\bar{\mathbb{Q}}^G$.*

Recall that for an elliptic curve E over a number field F, the *analytic rank* $\mathrm{rk}_{an}\,E/F$ is defined as $\mathrm{ord}_{s=1}L(E/F,s)$. Even for E defined over $\mathbb{Q}$ the analytic continuation of $L(E/F,s)$ to $s=1$ is not known for general F, but in the proof of the theorem we will choose number fields where it is. The *p^∞-Selmer rank* $\mathrm{rk}_p\,E/F$ is the usual Mordell-Weil rank $\mathrm{rk}\,E/F=\dim E(F)\otimes\mathbb{Q}$ plus the $\mathbb{Z}_p$-corank of $Ш(E/F)$, the number of copies of $\mathbb{Q}_p/\mathbb{Z}_p$ in $Ш(E/F)$. We remind the reader of the usual conjectures concerning these ranks:

Conjectures. *Let K be a number field, and E/K an elliptic curve.*

- $L(E/K,s)$ *is entire, and* $\mathrm{rk}\,E/K=\mathrm{rk}_{an}E/K$ *(Birch–Swinnerton-Dyer).*
- $Ш(E/K)$ *is finite; thus* $\mathrm{rk}\,E/K=\mathrm{rk}_p\,E/K$ *for all p (Shafarevich-Tate).*
- $(-1)^{\mathrm{rk}\,E/K}=w(E/K)$ *(Parity conjecture).*

The global root number $w(E/K)$ is defined as a product of local root numbers $w(E/K_v)=\pm1$ over all places of K, and is the conjectural sign in the

---

Date: March 6, 2008.

MSC 2000: Primary 11G05; Secondary 11G40, 14G25.

[†]Supported by a Royal Society University Research Fellowship.

functional equation for $L(E/K, s)$. Thus it determines $\mathrm{rk}_{an}(E/K) \mod 2$, so the parity conjecture is a weaker version of the Birch–Swinnerton-Dyer conjecture.

In fact, the proof of Theorem 2 shows that the parity conjecture also implies Conjecture 1 for elliptic curves over $K = \mathbb{Q}$. When K is a general number field, the analytic continuation of $L(E/K, s)$ to $s = 1$ is not known so we cannot say anything about $\mathrm{rk}_{an}\, E/K$. However, we still have the following

Theorem 3. *Let K be a number field, E/K an elliptic curve and G a finitely generated subgroup of $\mathrm{Gal}(\bar{K}/K)$. Then both the parity and the Shafarevich-Tate conjectures imply Conjecture 1, provided*

- *K has a real place, or*
- *E has non-integral j-invariant.*

Remark 4. Slightly more generally, the theorem applies if there is a place v of K and some quadratic extension L_w/K_v where $w(E/L_w) = -1$. (See [4] for a classification of such E/K_v.)

The proofs of both theorems rely on an elementary root number argument in $\mathbb{F}_p^r \rtimes C_2$-extensions and some version of the parity conjecture. For the second theorem we will need the following result, which is a slight variation of [3] Thm. 1.3.

Proposition 5. *Let K be a number field, E/K an elliptic curve and M/K a quadratic extension. Suppose for every prime $w|6$ of M which is not split in M/K, E has semistable reduction at w, and the reduction is not good supersingular if $w|2$. If $\text{Ш}(E/M(E[2]))[6^\infty]$ is finite, then*

$$(-1)^{\mathrm{rk}\, E/M} = w(E/M).$$

This has a consequence, which is of interest in itself:

Corollary 6. *Let M/K be a quadratic extension of number fields, E/K an elliptic curve and assume that $\text{Ш}(E/M(E[2]))$ is finite. If all primes of bad reduction of E split in M/K, then the parity conjecture holds for E/M,*

$$(-1)^{\mathrm{rk}\, E/M} = w(E/M) = (-1)^{\#\text{Archimedean places of } M}.$$

Notation. Throughout the note K denotes a number field and E an elliptic curve defined over K. We write $w(E/K_v) = \pm 1$ for the local root number of E at v, and $w(E/K) = \prod_v w(E/K_v)$ for the global root number. For an Artin representation τ of $\mathrm{Gal}(\bar{K}/K)$ we write $w(E/K_v, \tau)$ and $w(E/K, \tau) = \prod_v w(E/K_v, \tau)$ for the local and global root numbers of the twist of E by τ. See e.g. [10] for properties of root numbers of elliptic curves and their twists.

Acknowledgments. We would like to thank A. Jensen for helpful discussions.

2. Proof of Theorem 2

Lemma 7. *Let M/K be a quadratic extension of number fields, E/K an elliptic curve and $G \subset \mathrm{Gal}(\bar{K}/K)$ a finitely generated subgroup. For every odd prime p and $r \geq 1$ there exists a Galois extension F/K containing M such that*

1. $\mathrm{Gal}(F/K) \cong \mathbb{F}_p^r \rtimes C_2$*, with* C_2 *acting by* -1*.*
2. *The image of* G *in* $\mathrm{Gal}(F/K)$ *has order at most 2.*
3. *The primes of* M *above primes of bad reduction for* E/K *split completely in* F*.*

For such an extension, $w(E/K, \rho) = w(E/M)$ *for every irreducible 2-dimensional representation* ρ *of* $\mathrm{Gal}(F/K)$*.*

Proof. Pick primes $\wp_1, \ldots, \wp_n$ of K that split completely in $M(\zeta_p)$, and consider $\mathfrak{m} = \prod_i \wp_i$ as a modulus of M (ζ_p denotes a primitive pth root of unity). Write $I^{\mathfrak{m}}$ for the group of fractional ideals of M that are coprime to $\mathfrak{m}$, and $P^{\mathfrak{m}}$ for the subgroup of principal ideals that can be generated by an element congruent to 1 mod $\mathfrak{m}$.

Let Q be the largest $\mathbb{F}_p$-vector space quotient of the group $I^{\mathfrak{m}}/P_{\mathfrak{m}}$. Then $2n - \delta \leq \dim Q \leq 2n$ with $\delta = \dim \mathcal{O}_M^*/\mathcal{O}_M^{*p} + \dim \mathrm{Cl}_M[p]$. Comparing this with the corresponding group of K, we see that the $\mathrm{Gal}(M/K)$-antiinvariant part of Q has dimension $d \geq n - \delta$. By class field theory it yields a Galois extension F_n/K with $\mathrm{Gal}(F_n/K) \cong \mathbb{F}_p^d \rtimes C_2$, C_2 acting by -1.

Consider $H = \mathrm{Gal}(F_n/M) = \mathbb{F}_p^d$. The group $G \cap \mathrm{Gal}(\bar{M}/M)$ is of index at most 2 in G, and the $\mathbb{F}_p$-dimension of its image in H is bounded by the number of generators of G. Also, the decomposition subgroup in H of any prime of M has size bounded by a constant independent of n. Now let $H_0 < H$ be generated by the image of $G \cap \mathrm{Gal}(\bar{M}/M)$ and the decomposition subgroups of primes of M that lie above primes of bad reduction for E/K. Then $F_n^{H_0}$ satisfies (2) and (3), and letting $n \to \infty$ gives the desired extensions F/K.

For the last claim, let ϵ be the non-trivial character of $\mathrm{Gal}(M/K)$. The complex irreducible representations of $\mathrm{Gal}(F/K)$ are $\mathbf{1}$, ϵ and 2-dimensional representations each of which factors through a D_{2p}-quotient.

Recall that $w(E/M) = w(E/K, \mathbf{1} \oplus \epsilon)$ by inductivity of global root numbers, so it suffices to check that $w(E/K_v, \mathbf{1} \oplus \epsilon) = w(E/K_v, \rho)$ for all places v of K. This holds for places v whose decomposition group D in F/K has order at most 2, because ρ and $\mathbf{1} \oplus \epsilon$ are isomorphic as D-representations. In particular, this includes Archimedean places and places of bad reduction for E. On the other hand, if E has good reduction at v, then for every 2-dimensional self-dual Artin representation τ of $\mathrm{Gal}(\bar{K}/K)$,

$$w(E/K_v, \tau) = w(\tau/K_v)^2 = \det(\tau(-1)),$$

by the unramified twist formula [12] 3.4.6 and the determinant formula [12] 3.4.7. Here we implicitly use the local reciprocity map at v to evaluate τ at -1. As $\det \rho = \epsilon = \det(\mathbf{1} \oplus \epsilon)$, the claim follows. □

We now prove Theorem 2. Take an imaginary quadratic field $M=\mathbb{Q}(\sqrt{-d})$ where all bad primes for E split. Note that $w(E/M)=-1$, as the contribution from bad primes is $(\pm 1)^2$, and the local root number is $+1$ for primes of good reduction and -1 for infinite places.

Let p be an odd prime. By the p-parity conjecture for $E/\mathbb{Q}$ and its quadratic twist by $-d$ ([3] Thm. 1.4), we have $(-1)^{\mathrm{rk}_p(E/M)}=w(E/M)$, so $\mathrm{rk}_p(E/M)$ is odd. Let F be as in Lemma 7 for some $r\geq 1$, and let $V\subset\mathbb{F}_p^r$ be any index p subgroup, so $V\triangleleft\mathrm{Gal}(F/\mathbb{Q})$ and $\mathrm{Gal}(F^V/\mathbb{Q})\cong D_{2p}$. The image of G in this Galois group has order at most 2, so there is a degree p extension $L/\mathbb{Q}$ inside F^V fixed by G. As all bad primes for E split in M, by [3] Prop. 4.17,

$$\mathrm{rk}_p\, E/M+\tfrac{2}{p-1}(\mathrm{rk}_p\, E/L-\mathrm{rk}_p\, E/\mathbb{Q})\equiv 0 \pmod 2.$$

In particular, $\mathrm{rk}_p\, E/L>\mathrm{rk}_p\, E/\mathbb{Q}$.

Write X for the dual p^∞-Selmer group $\mathrm{Hom}(\mathrm{Sel}_{p^\infty}(E/F),\mathbb{Q}_p/\mathbb{Z}_p)$ and $\mathcal{X}=X\otimes_{\mathbb{Z}_p}\mathbb{Q}_p$. Thus $\mathcal{X}$ is a $\mathbb{Q}_p$-valued representation of $\mathrm{Gal}(F/\mathbb{Q})$, and $\mathrm{rk}_p\, E/k=\dim\mathcal{X}^{\mathrm{Gal}(F/k)}$ for every $k\subset F$, see e.g. [3] Lemma 4.14.

The irreducible $\mathbb{Q}_p$-representations of $\mathrm{Gal}(F^V/\mathbb{Q})\cong D_{2p}$ are trivial $\mathbf{1}$, sign ϵ and $(p-1)$-dimensional irreducible ρ_V. Their invariants under an element of order 2 of D_{2p} are 1-, 0- and $\frac{p-1}{2}$-dimensional, respectively. It follows that $\mathcal{X}$ contains a copy of ρ_V for every V, and $\dim\mathcal{X}^G$ is therefore at least $\frac{p-1}{2}$ times the number of hyperplanes $V\subset\mathbb{F}_p^r$ (which is $\frac{p^r-1}{p-1}$). Letting $r\to\infty$ we deduce that the p^∞-Selmer rank of E is unbounded for number fields inside $\bar{\mathbb{Q}}^G$.

To prove that the analytic rank is unbounded, let $p=3$ for simplicity and take the same F as above. The irreducible complex representations of $\mathrm{Gal}(F/\mathbb{Q})$ are $\mathbf{1},\epsilon$, and ρ_V for varying V. The curve $E/\mathbb{Q}$ is modular, so the L-functions $L(E/\mathbb{Q},s)$, $L(E,\epsilon,s)$ and $L(E,\rho_V,s)$ are analytic and satisfy the expected functional equation. (The twists by ρ_V are Rankin-Selberg products.) The same applies to $L(E/k,s)$ for every $k\subset F$; indeed, by Artin formalism,

$$L(E/k,s) \;=\; L(E,\mathrm{Ind}_k^{\mathbb{Q}}\mathbf{1},s) \;=\; \prod_{\tau\in\{\mathbf{1},\epsilon\}\cup\{\rho_V\}_V} L(E,\tau,s)^{\langle\tau,\mathrm{Ind}_k^{\mathbb{Q}}\mathbf{1}\rangle},$$

with $\mathrm{Ind}_k^{\mathbb{Q}}$ a shorthand for $\mathrm{Ind}_{\mathrm{Gal}(F/k)}^{\mathrm{Gal}(F/\mathbb{Q})}$. Finally, $w(E,\rho_V)=-1$ by Lemma 7, so each $L(E,\rho_V,s)$ vanishes at $s=1$. Hence

$$\mathrm{ord}_{s=1}L(E/k,s)\geq\frac{[k:\mathbb{Q}]-2}{2}$$

as $\mathrm{Ind}_k^{\mathbb{Q}}\mathbf{1}$ contains at most one copy of $\mathbf{1}$ and ϵ. Now take $k=F^G$ and let $r\to\infty$ as before.

3. Proof of Theorem 3

We claim there is a quadratic extension M/K where $w(E/M) = -1$ and $\operatorname{rk} E/M$ is odd. This will suffice, as we can then use exactly the same construction as in the proof of Theorem 2 with K in place of $\mathbb{Q}$ and any odd p.

Fix a place v of K which is either real or with $\operatorname{ord}_v j(E) < 0$. We take M/K to be any quadratic extension where

- v becomes complex if $K_v = \mathbb{R}$.
- There is a unique prime v' above v, and E has split multiplicative reduction at v' if $\operatorname{ord}_v j(E) < 0$.
- If $w \neq v$ is either Archimedean, a place of bad reduction of E or divides 2 then w splits in M/K.

Such a quadratic M exists by the weak approximation theorem, as we prescribe only its local behaviour at finitely many places. The existence of an appropriate local extension in the second case follows from the theory of the Tate curve, see e.g. [11] Thm 5.3.

Write v' for the unique prime of M above v. Then $w(E/M_{v'}) = -1$ (see e.g. [10] Thm. 2) and all other Archimedean places and bad primes for E contribute $(\pm 1)^2 = 1$, so $w(E/M) = -1$. Finally $\operatorname{rk} E/M$ is odd, by either the assumed parity conjecture or the Shafarevich-Tate conjecture together with Proposition 5.

4. Proof of Proposition 5

Lemma 8. *Let K be a number field and E/K an elliptic curve with a K-rational 2-torsion point. Let Σ be the set of primes $v|2$ of K where E has additive or good supersingular reduction. Suppose M/K is a quadratic extension where every prime $v \in \Sigma$ either splits or becomes a place of multiplicative or good ordinary reduction for E. Then $(-1)^{\operatorname{rk}_2 E/M} = w(E/M)$.*

Proof. Choose a model for E/K of the form

$$E: \ y^2 = x^3 + ax^2 + bx \ ,$$

and let $\phi : E \to E'$ be the 2-isogeny with $(0,0)$ in the kernel.

Let $\sigma_w(E/M) = (-1)^{\operatorname{ord}_2 \operatorname{coker} \phi_w - \operatorname{ord}_2 \ker \phi_w}$, where $\phi_w : E(M_w) \to E'(M_w)$ is the induced map on local points. By Cassels' formula (see [2] §1),

$$(-1)^{\operatorname{rk}_2 E/M} = \prod_w \sigma_w(E/M).$$

Write $(\cdot,\cdot)_w = \pm 1$ for the Hilbert symbol at w. Then

$$\prod_{w|v} w(E/M_w) = \prod_{w|v} \sigma_w(E/M)(a,-b)_w(-2a, a^2-4b)_w$$

for every place v of K. Indeed, this trivially holds for all v that split in M/K, and it holds at all other primes by [2], proof of Thm. 4 in §7. The result follows by the product formula for Hilbert symbols. □

Lemma 9. *Let K be a number field and E/K an elliptic curve. Let Σ be the set of primes $v|6$ of K where E has additive reduction. Suppose M/K is a quadratic extension where every prime $v \in \Sigma$ either splits or becomes a place of semistable reduction for E. Let F/K be any S_3-extension not containing M, and write $N = F^{C_2}M$, $N' = F^{C_3}M$. Then*

$$w(E/M)w(E/N)w(E/N') = (-1)^{\mathrm{rk}_3 E/M + \mathrm{rk}_3 E/N + \mathrm{rk}_3 E/N'}.$$

Proof. Fix a global differential $\omega \neq 0$ for E/K. For an extension k/K and a prime w of k write $C_w(E/k) = c_w(E/k)|\omega/\omega_w^o|_w$, where c_w is the local Tamagawa number, ω_w^o a Néron differential at w, and $|\cdot|_w$ the normalised w-adic absolute value. By [3] Thm. 4.11 with $p = 3$,

$$\mathrm{rk}_3 E/M + \mathrm{rk}_3 E/N + \mathrm{rk}_3 E/N' \equiv \mathrm{ord}_3 \frac{\prod_z C_z(E/MF)}{\prod_u C_u(E/N')} \mod 2,$$

the products taken over the primes of MF and N' respectively. It suffices to show that for every place v of K,

$$\prod_{s|v} w(E/M_s) \prod_{t|v} w(E/N_t) \prod_{u|v} w(E/N'_u) = \frac{\prod_{z|v}(-1)^{\mathrm{ord}_3 C_z(E/MF)}}{\prod_{u|v}(-1)^{\mathrm{ord}_3 C_u(E/N')}},$$

where we interpret the right-hand side as 1 for Archimedean places. If v splits in M/K the formula trivially holds as each term occurs an even number of times. For all other v, this is proved in [3] Prop 3.3 (the G-set argument in Case 1 also covers Archimedean places). □

We now prove Proposition 5.

The assumption on Ш also forces Ш$(E/k)[6^\infty]$ to be finite for all intermediate fields $K \subset k \subset M(E[2])$, see e.g. [3] Rmk. 2.10. Now we proceed as in the proof of [3] Thm. 3.6.

Write $F = K(E[2])$. If $E(K)[2] \neq 0$, apply Lemma 8. Otherwise $\mathrm{Gal}(F/K)$ is either C_3 or S_3. In the former case, $\mathrm{Gal}(FM/M) \cong C_3$ as well, thus $\mathrm{rk}\, E/M$ and $\mathrm{rk}\, E/FM$ have the same parity. It is also well-known that global root numbers are unchanged in odd degree cyclic extensions (this follows from [12] 3.4.7, 4.2.4), so $w(E/FM) = w(E/M)$ and the result again follows from Lemma 8 applied to FM/F.

In the last case $\mathrm{Gal}(F/K) \cong S_3$, write $N = F^{C_2}M$, $N' = F^{C_3}M$. If $M \not\subset F$, the above argument shows that $w(E/N) = (-1)^{\mathrm{rk}\, E/N}$ and similarly for N'; now apply Lemma 9. If $M \subset F$, then F/M is a Galois cubic extension, so we may again show that $w(E/F) = (-1)^{\mathrm{rk}\, E/F}$. But this holds by Lemma 8 applied to the quadratic extension F/N.

Finally, Corollary 6 is immediate from the fact that $w(E/K_v) = -1$ for Archimedean v.

References

[1] F. Breuer, B.-H. Im, Heegner points and the rank of elliptic curves over large extensions of global fields, arxiv:math/0604107.

[2] T. Dokchitser, V. Dokchitser, Parity of ranks for elliptic curves with a cyclic isogeny, J. Number Theory 128 (2008), 662–679.

[3] T. Dokchitser, V. Dokchitser, On the Birch–Swinnerton-Dyer quotients modulo squares, 2006, arxiv: math.NT/0610290.

[4] T. Dokchitser, V. Dokchitser, Elliptic curves with all quadratic twists of positive rank, 2008, arxiv: 0802.4027.

[5] B.-H. Im, Heegner points and Mordell-Weil groups of elliptic curves over large fields, to appear in Trans. Amer. Math. Soc.

[6] B.-H. Im, Mordell-Weil groups and the rank of elliptic curves over large fields, arxiv:math/0411533.

[7] B.-H. Im, Heegner points and Mordell-Weil groups of elliptic curves over large fields, arxiv:math/0411534.

[8] M. Larsen, Rank of elliptic curves over almost algebraically closed fields, Bull. London Math. Soc. 35 (2003), 817-820.

[9] Á. Lozano-Robledo, On the product of twists of rank two and a conjecture of Larsen, preprint.

[10] D. Rohrlich, Galois Theory, elliptic curves, and root numbers, Compos. Math. 100 (1996), 311–349.

[11] J. H. Silverman, Advanced Topics in the Arithmetic of Elliptic Curves, GTM 151, Springer-Verlag 1994.

[12] J. Tate, Number theoretic background, in: Automorphic forms, representations and L-functions, Part 2 (ed. A. Borel and W. Casselman), Proc. Symp. in Pure Math. 33 (AMS, Providence, RI, 1979) 3-26.

Robinson College, Cambridge CB3 9AN, United Kingdom
E-mail address: `t.dokchitser@dpmms.cam.ac.uk`

Gonville & Caius College, Cambridge CB2 1TA, United Kingdom
E-mail address: `v.dokchitser@dpmms.cam.ac.uk`